

# PATIENT CONSENT FORM

## FOR SEASONAL INFLUENZA VACCINE

I have read, or have had explained to me, the CDC Vaccine Information Statement about influenza and the influenza vaccine. I understand that this vaccine may cause flu-like symptoms in some people and in rare incidents Guillain-Barré syndrome. I have had an opportunity to ask questions which were answered to my satisfaction. I understand the benefits and risks of influenza vaccine and request that the vaccine be given to me (or person named below for whom I am authorized to make this request).

Please print:

Name: \_\_\_\_\_ Date of Birth: //  
(FIRST) (MIDDLE) (LAST)

Parent or Guardian's Name (if applicable): \_\_\_\_\_

Has the person receiving the vaccine ever had a severe allergic (hypersensitivity) reaction to eggs, chickens, or chicken feathers? \_\_\_Yes \_\_\_No

Does the person receiving the vaccine have a history of Guillain-Barré syndrome or a persistent neurological illness? \_\_\_Yes \_\_\_No

Is the person receiving the vaccine pregnant? \_\_\_Yes \_\_\_No (If yes, LAIV contraindicated, TIV recommended)

Is the person receiving the vaccine allergic to Thimerosal (Preservative found in contact lens solution), any vaccine ingredient, or latex? \_\_\_Yes \_\_\_No

For child 6 mo-8 yrs, have they received 2 or more doses of influenza vaccine since July 2010? \_\_\_Yes \_\_\_No  
(If no, the child will need to receive 2 vaccinations [at least one month apart] for the best protection against flu.)

\_\_\_\_\_  
Signature of person receiving vaccine OR Parent/Guardian

\_\_\_\_\_  
Date

**DO NOT WRITE IN THIS SPACE—OFFICE USE ONLY VIS Edition Provided:** \_\_\_\_\_

Lot number: \_\_\_\_\_ Expiration Date: \_\_\_\_\_

Dose #1 or  Dose #2  
(Circle One - Pediatric Only)

**LAIV Nasal spray is recommended for children aged 2-8 (older adolescents and adults may receive as well if stock allows).**

CHECK ONE:

- \_\_\_ 0.5 mL IM Influenza Virus Vaccine given in \_\_\_left \_\_\_right deltoid –  TIV or  QIV  
\_\_\_ 0.5 mL IM Influenza HIGH Dose Virus Vaccine given in \_\_\_left \_\_\_right deltoid (65+) TIV-SR  
\_\_\_ 0.2 mL Live Attenuated Influenza Virus Vaccine given intranasally (half each nostril)  
\_\_\_ 0.5mL FluBlok Influenza Virus Vaccine given in \_\_\_left \_\_\_right deltoid  
\_\_\_ Children 6-35 months: 0.25 mL/dose given in \_\_\_left \_\_\_right deltoid (1 or 2 doses per season)  
\_\_\_ Children 3-8 years: 0.5 mL/dose given in \_\_\_left \_\_\_right deltoid (1 or 2 doses per season)  
\_\_\_ Children older than 9 years: 0.5 mL/dose given in \_\_\_left \_\_\_right deltoid (1 dose per season)

\_\_\_\_\_  
Nurse/MA/Provider's Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Time

